

Singular Loci of Varieties of Complexes, II

Nicolae Gongiulea

Department of Mathematics, University of Michigan, Ann Arbor, Michigan 48109

E-mail: ngongiul@math.lsa.umich.edu

Communicated by Peter Littelmann

Received November 20, 1998

In this paper we determine the singular loci of the irreducible components of the variety of complexes. © 2001 Academic Press

INTRODUCTION

Let W_1, \dots, W_{h+1} be finite-dimensional vector spaces over some field k such that $\dim W_i = n_i$ for $1 \leq i \leq h+1$. Let Z be the affine space of all h -tuples of linear maps

$$(f_1, \dots, f_h): W_1 \xrightarrow{f_1} W_2 \xrightarrow{f_2} \dots \xrightarrow{f_{h-1}} W_h \xrightarrow{f_h} W_{h+1}.$$

The subvariety \mathcal{E} of Z defined by

$$\mathcal{E} = \{(f_1, f_2, \dots, f_h) \in Z \mid f_{i+1}f_i = 0 \text{ for } 1 \leq i \leq h-1\}$$

is called the *variety of complexes*.

For positive integers k_1, \dots, k_h such that $k_i + k_{i+1} \leq n_{i+1}$ for $0 \leq i \leq h$ (where $k_0 = k_{h+1} = 0$), the varieties

$$V(k_1, \dots, k_h) = \{(f_1, \dots, f_h) \in \mathcal{E} \mid \text{rank}(f_i) \leq k_i \text{ for } 1 \leq i \leq h\}$$

are irreducible subvarieties of \mathcal{E} .

In this paper we determine (the irreducible components of) the singular locus of $V(k_1, \dots, k_h)$. Let $V = V(k_1, \dots, k_h)$,

$$V_i = V(k_1, \dots, k_{i-1}, k_i - 1, k_{i+1}, \dots, k_h) \quad \text{for } 1 \leq i \leq h,$$

$$V_{j,j+1} = V(k_1, \dots, k_{j-1}, k_j - 1, k_{j+1} - 1, k_{j+2}, \dots, k_h)$$

$$\text{for } 1 \leq j \leq h-1.$$

We show the following

THEOREM 1. *The irreducible components of $\text{Sing } V$ are V_i , with $i \in \Omega$, and $V_{j,j+1}$, with $j \notin \Omega$, $j+1 \notin \Omega$, where Ω is the set of all $1 \leq i \leq h$ such that $k_{i-1} + k_i < n_i$ and $k_i + k_{i+1} < n_{i+1}$.*

As a consequence, we obtain

THEOREM 2. *If $V = V(k_1, \dots, k_h)$ is an irreducible component of the variety of complexes \mathcal{C} , then the irreducible components of $\text{Sing } V$ are $V_{1,2}, \dots, V_{h-1,h}$.*

For $h = 2$, these results are proved in [3].

The varieties $V(k_1, \dots, k_h)$ can be identified with opposite cells of certain Schubert varieties (see [4, 5]); thus, using Theorem 1, one can determine the irreducible components of these Schubert varieties.

We thank V. Lakshmibai for helpful discussions during the preparation of this paper.

1. PRELIMINARIES

Let us fix the positive integers $h \geq 2$, n_1, \dots, n_{h+1} , and let $\mathbf{n} = (n_1, \dots, n_{h+1})$. Let W_1, \dots, W_{h+1} be finite-dimensional vector spaces over a field k with $\dim W_1 = n_1, \dots, \dim W_{h+1} = n_{h+1}$. Let Z be the affine space of all h -tuples of linear maps

$$(f_1, \dots, f_h): W_1 \xrightarrow{f_1} W_2 \xrightarrow{f_2} \dots \xrightarrow{f_{h-1}} W_h \xrightarrow{f_h} W_{h+1}.$$

The group $G_{\mathbf{n}} = GL(n_1) \times \dots \times GL(n_{h+1})$ acts on Z by

$$(g_1, g_2, \dots, g_{h+1}) \cdot (f_1, f_2, \dots, f_h) = (g_2 f_1 g_1^{-1}, g_3 f_2 g_2^{-1}, \dots, g_{h+1} f_h g_h^{-1}).$$

The subvariety \mathcal{C} of Z defined by

$$\mathcal{C} = \{(f_1, f_2, \dots, f_h) \in Z \mid f_{i+1} f_i = 0 \text{ for } 1 \leq i \leq h-1\}$$

is called the *variety of complexes*.

Let $\mathcal{K}_{\mathbf{n}}$ denote the set of all $\mathbf{k} = (k_1, \dots, k_h) \in \mathbb{Z}_+^h$, such that $k_i + k_{i+1} \leq n_{i+1}$ for $0 \leq i \leq h$ (here $k_0 = k_{h+1} = 0$). For $\mathbf{k} = (k_1, \dots, k_h) \in \mathcal{K}_{\mathbf{n}}$, let

$$V(\mathbf{k}) = \{(f_1, \dots, f_h) \in \mathcal{C} \mid \text{rank}(f_i) \leq k_i \text{ for } 1 \leq i \leq h\}.$$

Let X_i be the $n_{i+1} \times n_i$ matrix of indeterminates, for $1 \leq i \leq h$, and let $k[X_1, \dots, X_h]$ be the polynomial ring over the indeterminates in $X_1 \cup \dots \cup X_h$. For $\mathbf{k} \in \mathcal{K}_{\mathbf{n}}$, let $I(\mathbf{k})$ be the ideal in $k[X_1, \dots, X_h]$ generated by the entries of $X_{i+1} X_i$, for $1 \leq i \leq h-1$, and the $k_i + 1$ minors of X_i , for $1 \leq i \leq h$.

Let $G = GL(n)$. Let T be the maximal torus consisting of diagonal matrices in G , let B be the Borel subgroup consisting of the upper triangular matrices in G , and let $W = \mathcal{S}_n$ be the Weyl group of G . Let P_d , $1 \leq d \leq n-1$, be the maximal parabolic subgroups in G , where

$$P_d = \left\{ A \in G \mid A = \begin{pmatrix} * & * \\ 0_{(n-d) \times d} & * \end{pmatrix} \right\}.$$

Consider the parabolic subgroup $Q = P_{a_1} \cap \cdots \cap P_{a_h}$ of G , i.e., the subgroup consisting of all elements of the form

$$\begin{pmatrix} A_1 & * & * & \cdots & * & * \\ 0 & A_2 & * & \cdots & * & * \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & A_h & * \\ 0 & 0 & 0 & \cdots & 0 & A_{h+1} \end{pmatrix},$$

where A_i is a matrix of size $(a_i - a_{i-1}) \times (a_i - a_{i-1})$ for $1 \leq i \leq h+1$ (here $a_0 = 0$ and $a_{h+1} = n$). The Weyl group of Q is $W_Q = \mathcal{S}_{a_1} \times \mathcal{S}_{a_2 - a_1} \times \cdots \times \mathcal{S}_{n - a_h}$. The set of T -fixed points in G/Q for the action given by multiplication is precisely the set $\{e_{w,Q}, w \in W\}$, where for $w \in W$, $e_{w,Q}$ is the point in G/Q corresponding to the coset wQ . For $w \in W$, let $X_Q(w)$ be the *Schubert variety* in G/Q associated to wW_Q , i.e., the Zariski closure of $Be_{w,Q}$ in G/Q . It is well known that Schubert varieties are irreducible.

Let O^- be the set of elements of G of the form

$$\begin{pmatrix} I_1 & 0 & 0 & \cdots & 0 & 0 \\ * & I_2 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ * & * & * & \cdots & I_h & 0 \\ * & * & * & \cdots & * & I_{h+1} \end{pmatrix},$$

where I_j is the identity matrix of size $(a_j - a_{j-1}) \times (a_j - a_{j-1})$ for $1 \leq j \leq h+1$ (here $a_0 = 0$ and $a_{h+1} = n$). Then the restriction of the canonical morphism $G \rightarrow G/Q$ to O^- is an open immersion, and the image of O^- is isomorphic to the *opposite big cell* in G/Q , i.e., with $B^-e_{\text{id},Q}$, where B^- is the Borel subgroup opposite to B , consisting of the lower triangular matrices in G . Thus the set O^- is identified with the opposite big cell in G/Q . Any Schubert variety $X_Q(w) \subset G/Q$ has a non-empty intersection with the opposite big cell O^- , and $X_Q(w) \cap O^-$ is called the *opposite cell* in $X_Q(w)$.

The following result is proved in [5]:

THEOREM 1.1. *For $\mathbf{k} \in \mathcal{K}_{\mathbf{n}}$, the variety $V(\mathbf{k})$ is isomorphic to the opposite cell in a certain Schubert variety in $GL(n)/Q$, where $n = n_1 + \cdots + n_{h+1}$ and $Q = P_{a_1} \cap \cdots \cap P_{a_h}$, $a_i = n_1 + \cdots + n_i$ for $1 \leq i \leq h$. Furthermore, $I(\mathbf{k})$ defines the reduced scheme structure on $V(\mathbf{k})$.*

This isomorphism is the restriction of the map

$$\varphi: M(n_2 \times n_1) \times \cdots \times M(n_{h+1} \times n_h) \rightarrow O^-,$$

$$(A_1, \dots, A_h) \mapsto \begin{pmatrix} I_1 & 0 & 0 & \cdots & 0 & 0 & 0 \\ A_1 & I_2 & 0 & \cdots & 0 & 0 & 0 \\ 0 & A_2 & I_3 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & I_{h-1} & 0 & 0 \\ 0 & 0 & 0 & \cdots & A_{h-1} & I_h & 0 \\ 0 & 0 & 0 & \cdots & 0 & A_h & I_{h+1} \end{pmatrix} \pmod{Q}$$

to $V(\mathbf{k})$.

This result can also be obtained from the identification of *quiver varieties* (which are more general than the varieties $V(\mathbf{k})$ with $\mathbf{k} \in \mathcal{K}_{\mathbf{n}}$, discussed here) with opposite cells of Schubert varieties (see [4]).

It is easily seen that $V(\mathbf{k})$, with $\mathbf{k} = (k_1, \dots, k_h) \in \mathcal{K}_{\mathbf{n}}$, are all the closed irreducible $G_{\mathbf{n}}$ -stable subvarieties of \mathcal{E} and that $\mathcal{E} = \bigcup_{\mathbf{k} \in \mathcal{K}_{\mathbf{n}}} V(\mathbf{k})$.

Define the following partial order on the set $\mathcal{K}_{\mathbf{n}}$: for $\mathbf{k} = (k_1, k_2, \dots, k_h)$, $\mathbf{k}' = (k'_1, k'_2, \dots, k'_h)$ in $\mathcal{K}_{\mathbf{n}}$, $\mathbf{k} \geq \mathbf{k}' \Leftrightarrow k_i \geq k'_i$, $1 \leq i \leq h$. Clearly, $V(\mathbf{k}') \subseteq V(\mathbf{k})$ if and only if $\mathbf{k}' \leq \mathbf{k}$. Hence we have the following

THEOREM 1.2. *The irreducible components of \mathcal{E} are its subvarieties of the form $V(\mathbf{k})$, with \mathbf{k} a maximal element of $\mathcal{K}_{\mathbf{n}}$ (with respect to the partial order above).*

The following result is also proved in [5]:

THEOREM 1.3. *Let $\mathbf{k} \in \mathcal{K}_{\mathbf{n}}$.*

1. $\dim V(\mathbf{k}) = \sum_{1 \leq i \leq h+1} (n_i - k_i)(k_{i-1} + k_i)$, where $k_0 = k_{h+1} = 0$.
2. $\text{codim}_Z V(\mathbf{k}) = \sum_{i=1}^h (n_{i+1} - k_i)(n_i - k_i) + \sum_{i=1}^{h-1} k_i k_{i+1}$.

This theorem can also be deduced from [1].

Considering some fixed basis in each W_i , we have the identifications $W_i \cong k^{n_i}$ and $Z \cong M(n_2 \times n_1) \times \cdots \times M(n_{h+1} \times n_h)$, where $M(l \times m)$ denotes the affine space of matrices over k with l rows and m columns. Then \mathcal{E} can be identified with the set of all points (A_1, \dots, A_h) in

$M(n_2 \times n_1) \times \cdots \times M(n_{h+1} \times n_h)$ such that $A_2 A_1 = 0, \dots, A_h A_{h-1} = 0$. For $\mathbf{k} \in \mathcal{K}_n$, the variety $V(\mathbf{k})$ is identified with the set of all points (A_1, \dots, A_h) in $M(n_2 \times n_1) \times \cdots \times M(n_{h+1} \times n_h)$ such that $A_2 A_1 = 0, \dots, A_h A_{h-1} = 0$, $\text{rank } A_1 \leq k_1, \dots, \text{rank } A_h \leq k_h$. Note that $V(\mathbf{k})$ is the closure of the G_n -orbit in $M(n_2 \times n_1) \times \cdots \times M(n_{h+1} \times n_h)$ through a point (A_1, \dots, A_h) such that $\text{rank } A_1 = k_1, \dots, \text{rank } A_h = k_h$, and because of the action of G_n , we may assume, without loss of generality, that

$$A_i = \begin{pmatrix} 0 & 0 \\ M_i & 0 \end{pmatrix}, \quad \text{for } 1 \leq i \leq h,$$

where M_i is the $k_i \times k_i$ identity matrix.

2. SINGULAR LOCUS OF $V(k_1, \dots, k_h)$

In this section we determine the singular points of $V(\mathbf{k})$, using the Jacobian criterion. Recall that, by Theorem 1.1, $I(\mathbf{k})$ is the defining ideal of $V(\mathbf{k})$.

Let $\mathbf{k} = (k_1, \dots, k_h) \in \mathcal{K}_n$ and $V = V(k_1, k_2, \dots, k_h)$. For $1 \leq i \leq h$, let X_i denote the $n_{i+1} \times n_i$ matrix of variables, and let \mathcal{M}_i be the set of all $k_i + 1$ minors of X_i . Then V is the set of zeros of the polynomials in $\mathcal{M}_1, \dots, \mathcal{M}_h, X_2 X_1, \dots, X_h X_{h-1}$. Let $x = (A_1, \dots, A_h)$ be a point in V , A_i being an $n_{i+1} \times n_i$ matrix of rank at most k_i , and let J_x be the Jacobian matrix of V evaluated at x . The rows of J_x are indexed by the polynomials in the sets $\mathcal{M}_1, \dots, \mathcal{M}_h, X_2 X_1, \dots, X_h X_{h-1}$, while the columns of J are indexed by the entries in X_1, \dots, X_h . The (M, α) th entry of J_x , where M is a minor in $\mathcal{M}_1 \cup \cdots \cup \mathcal{M}_h$ and α is an entry in $X_1 \cup \cdots \cup X_h$, is equal to \pm the minor obtained from M by deleting the row and the column of α if α appears in M , and 0 otherwise. Let $f \in X_{i+1} X_i$ be the product of the r th row of X_{i+1} and the s th column of X_i , for some $1 \leq i \leq h-1$, $1 \leq r \leq n_{i+2}$, $1 \leq s \leq n_i$, and let α be an entry in $X_1 \cup \cdots \cup X_h$; then the (f, α) th entry of J_x is equal to the (j, s) th entry of X_i if α is the (r, j) th entry of X_i for some $1 \leq j \leq n_i$, the (r, l) th entry of X_{i+1} if α is the (l, s) th of X_i for some $1 \leq l \leq n_i$, and 0 otherwise.

The next lemma is obvious from the fact that the closure of the orbit through a point $x = (A_1, \dots, A_h)$ in V such that $\text{rank } A_i = k_i$ for all $1 \leq i \leq h$ is V itself. We include the proof of this result to establish notations that will be used in the following, and for improved understanding of the Jacobian matrix of V .

LEMMA 2.1. *Let $x = (A_1, \dots, A_h)$ be a point in V such that $\text{rank } A_i = k_i$ for all $1 \leq i \leq h$. Then x is a smooth point of V .*

Proof. Without loss of generality, we may assume that

$$A_i = \begin{pmatrix} 0 & 0 \\ M_i & 0 \end{pmatrix}, \quad \text{for } 1 \leq i \leq h,$$

where M_i is the $k_i \times k_i$ identity matrix.

Let us denote J_x by just J .

Next we describe all the nonzero rows of J .

For $1 \leq i \leq h$, a row of J indexed by a minor $M \in \mathcal{M}_i$ of X_i is nonzero if and only if M contains the identity block M_i . The only nonzero element in such a row is equal to ± 1 , placed in the column indexed by the upper right corner entry of M . When computing the rank of J , we may assume that such a ± 1 entry of J_x is actually equal to 1. Let \mathcal{R}_i be the set of all minors in \mathcal{M}_i containing M_i . Let Y_i be the set of all right upper corners of minors in \mathcal{R}_i , i.e., the set of all entries of X_i not contained in any of the rows or columns of M_i . Note that Y_i consists of the entries of the $(n_{i+1} - k_i) \times (n_i - k_i)$ right upper block of X_i , and hence its cardinality is $(n_{i+1} - k_i)(n_i - k_i)$. Also, $|\mathcal{R}_i| = (n_{i+1} - k_i)(n_i - k_i)$.

Now let f be an entry in $X_{i+1}X_i$ for some $1 \leq i \leq h - 1$. In Figure 1, the blocks X_i , X_{i+1} , $X_{i+1}X_i$ are displayed as shown below:

X_i	
$X_{i+1}X_i$	X_{i+1}

Let f be the product of the r th row of X_{i+1} and the s th column of X_i , $1 \leq r \leq n_{i+2}$, $1 \leq s \leq n_i$. Then the row of J indexed by f is nonzero if and only if either the r th row of X_{i+1} is nonzero (i.e., it is one of the last k_{i+1} rows of X_{i+1}) or the s th column of X_i is nonzero (i.e., it is one of the first k_i columns of X_i). The set of all such f 's in $X_{i+1}X_i$ can be written as $B_{i,i+1} \cup L_{i,i+1}$, where $B_{i,i+1}$ denotes the set of all entries in the last k_{i+1} rows of $X_{i+1}X_i$, and $L_{i,i+1}$ denotes the set of all entries in the first k_i columns of $X_{i+1}X_i$.

Now we find a maximal set of linearly independent rows of J among the set of rows described above.

The rows of J indexed by minors in $\mathcal{R}_1, \dots, \mathcal{R}_h$ are linearly independent, since their nonzero entries are in different columns (these columns are indexed by the entries in Y_1, \dots, Y_h , and each of them contains precisely one nonzero entry in the rows indexed by $\mathcal{R}_1, \dots, \mathcal{R}_h$).

For $1 \leq i \leq h - 1$, let $K_{i,i+1} = B_{i,i+1} \cap L_{i,i+1}$ (note that $K_{i,i+1}$ is the $k_{i+1} \times k_i$ left lower block of $X_{i+1}X_i$). A row of J indexed by $f \in K_{i,i+1}$ contains precisely two nonzero entries (equal to 1), neither of which is in




FIGURE 1

the same column with a nonzero entry of another nonzero row of J . The set of nonzero entries in the rows indexed by all $f \in K_{i,i+1}$ is in the columns indexed by the entries in the left upper block of X_i of size $k_{i+1} \times k_i$, which we denote by $Q_{i,i+1}^+$, and the right lower block of X_{i+1} of size $k_{i+1} \times k_i$, which we denote by $Q_{i,i+1}^-$.

A row of J indexed by $f \in (B_{i,i+1} \cup L_{i,i+1}) \setminus K_{i,i+1}$ contains precisely one nonzero entry (equal to 1) in a column indexed either by an element in the first k_{i+1} rows of Y_i or by an element in the last k_i columns of Y_{i+1} . Therefore these rows are identical to some of the rows indexed by minors in $\mathcal{R}_1, \dots, \mathcal{R}_h$.

Consequently, the rows of J indexed by minors in $\mathcal{R}_1, \dots, \mathcal{R}_h$ and entries of $K_{1,2}, \dots, K_{h-1,h}$ give a maximal set of linearly independent rows of J . Therefore,

$$\begin{aligned} \text{rank } J &= |\mathcal{R}_1| + \dots + |\mathcal{R}_h| + |K_{1,2}| + \dots + |K_{h-1,h}| \\ &= (n_1 - k_1)(n_2 - k_2) + \dots + (n_h - k_h)(n_{h+1} - k_h) \\ &\quad + k_1 k_2 + \dots + k_{h-1} k_h \\ &= \text{codim } V. \end{aligned}$$

This shows that x is a smooth point of V . ■

Notation. For the rest of the paper,

$$J, Y_1, \dots, Y_h, Q_{1,2}^+, Q_{1,2}^-, \dots, Q_{h-1,h}^+, Q_{h-1,h}^-$$

will denote the Jacobian matrix, respectively the sets, associated, as in Lemma 2.1, to the point (A_1, \dots, A_h) in V with

$$A_i = \begin{pmatrix} 0 & 0 \\ M_i & 0 \end{pmatrix}, \quad \text{for } 1 \leq i \leq h,$$

where M_i is the $k_i \times k_i$ identity matrix.

Remark 2.2. Note that the nonzero columns of J are the columns indexed by the entries in $Y_1, \dots, Y_h, Q_{1,2}^+, Q_{1,2}^-, \dots, Q_{h-1,h}^+, Q_{h-1,h}^-$. For $1 \leq j \leq h-1$, each column indexed by an entry in $Q_{j,j+1}^+$ is identical to a column indexed by the corresponding entry in $Q_{j,j+1}^-$. A maximal set of linearly independent columns of J consists of all distinct nonzero columns of J . We have $\text{codim } V = |Y_1| + \dots + |Y_h| + |Q_{1,2}^+| + \dots + |Q_{h-1,h}^+|$ ($|Q_{j,j+1}^\pm|$ denotes the cardinality of each of $Q_{j,j+1}^+$ and $Q_{j,j+1}^-$). A nonzero column whose index is not included in the set of indices of the columns in a maximal set Σ of linearly independent columns of J is indexed by an entry in $Q_{j,j+1}^+$ or $Q_{j,j+1}^-$ for some j , and it is identical to a column whose index belongs to $Q_{j,j+1}^-$ respectively $Q_{j,j+1}^+$ and is among the indices of columns in Σ .

Remark 2.3. For $1 \leq j \leq h$, a maximal set of linearly independent columns of J indexed by entries in $X_1, \dots, X_{j-1}, X_{j+1}, \dots, X_h$ consists of all such distinct nonzero columns. The cardinality of such a maximal set is $\text{codim } V - |Y_j|$. The set of indices of the columns in such a maximal set must include $Q_{j-1,j}^+$ and $Q_{j,j+1}^-$.

Remark 2.4. For $1 \leq j \leq h-1$, a maximal set of linearly independent columns of J indexed by $X_1, \dots, X_{j-1}, X_{j+2}, \dots, X_h$ consists of all such distinct nonzero. The cardinality of such a maximal set is $\text{codim } V - (|Y_j| + |Y_{j+1}| + |Q_{j,j+1}^\pm|)$. The set of indices of the columns in such a maximal set must include $Q_{j-1,j}^+$ and $Q_{j+1,j+2}^-$.

LEMMA 2.5. Let $x = (A_1, \dots, A_h)$ be a point of V such that $\text{rank } A_j < k_j$ for some j , $1 \leq j \leq h$. If x is smooth, then either $k_{j-1} + k_j = n_j$ or $k_j + k_{j+1} = n_{j+1}$ (here $k_0 = k_{h+1} = 0$).

Proof. Let $t_i = \text{rank } A_i$, for $1 \leq i \leq h$, with $t_j < k_j$. Without loss of generality, we may assume that

$$A_i = \begin{pmatrix} 0 & 0 \\ N_i & 0 \end{pmatrix}, \quad \text{for } 1 \leq i \leq h,$$

where N_i is the $t_i \times t_i$ identity matrix. As noted in Remark 2.3, there are precisely $\text{codim } V - |Y_j|$ linearly independent columns of J indexed by entries in $X_1, \dots, X_{j-1}, X_{j+1}, \dots, X_h$. Since the nonzero columns of J_x are

among the nonzero columns of J , the set T of linearly independent columns of J_x indexed by entries in $X_1, \dots, X_{j-1}, X_{j+1}, \dots, X_h$ has cardinality at most $\text{codim } V - |Y_j|$ (note that T consists of all distinct nonzero columns of J_x). Note that a column of J indexed by an entry in $X_1, \dots, X_{j-1}, X_{j+1}, \dots, X_h$ is either identical to some column indexed by an entry in X_j or it is linearly independent of the columns indexed by entries in X_j . Since x is smooth, $\text{rank } J_x = \text{codim } V$, so there exists a set S of cardinality at least $|Y_j|$ of linearly independent columns of J_x indexed by entries in X_j such that $S \cup T$ is a set of linearly independent columns of J_x . The only nonzero columns indexed by entries in X_j are actually indexed by entries in the first k_{j+1} rows of X_j and the last k_{j-1} columns of X_j (since $\text{rank } A_j < k_j$). The columns indexed by entries in $Q_{j-1,j}^- \cup Q_{j,j+1}^+$ cannot be in S , since they are in T (see Remark 2.3), and $S \cup T$ is a set of linearly independent columns of J_x . Thus S consists of columns indexed by entries in the first k_{j+1} rows of Y_j and the last k_{j-1} columns of Y_j . Since the cardinality of S must be at least $|Y_j|$, we deduce that the set of entries in the first k_{j+1} rows and the last k_{j-1} columns of Y_j must actually contain the set of entries in Y_j . This implies that either $k_{j-1} + k_j = n_j$ or $k_j + k_{j+1} = n_{j+1}$. ■

LEMMA 2.6. *Let $x = (A_1, \dots, A_h)$ be a point of V such that $\text{rank } A_j < k_j$ for some j , $1 \leq j \leq h$, and $\text{rank } A_i = k_i$ for $1 \leq i \leq h$, $i \neq j$. Then x is smooth if and only if either $k_{j-1} + k_j = n_j$ or $k_j + k_{j+1} = n_{j+1}$ (here $k_0 = k_{h+1} = 0$).*

Proof. In view of Lemma 2.5, we only have to show that if $k_{j-1} + k_j = n_j$ or $k_j + k_{j+1} = n_{j+1}$ then x is smooth. It is easily seen that the columns of J_x that are indexed by the elements in the disjoint sets

$$Y_1, \dots, Y_{j-1}, Y_{j+1}, \dots, Y_h, Q_{1,2}^-, \dots, Q_{j-2,j-1}^-, Q_{j+1,j+2}^+, \dots, Q_{h-1,h}^+, \Gamma_j,$$

where Γ_j is the set of entries in the first k_{j+1} rows and the last k_{j-1} columns of X_j , are linearly independent. Note that Y_j is contained in Γ_j (since $k_{j-1} + k_j = n_j$ or $k_j + k_{j+1} = n_{j+1}$), and the cardinality of Γ_j is precisely $|Y_j| + |Q_{j-1,j}^-| + |Q_{j,j+1}^+|$. Since the number of these columns is $\text{codim } V$, we conclude that $\text{rank } J_x = \text{codim } V$, and hence x is smooth. ■

Remark 2.7. An alternate proof of Lemma 2.6 can be obtained as follows. First note that if $k_{j-1} + k_j = n_j$, then $|L_{j-1,j}| = |\mathcal{R}_j|$, and if $k_j + k_{j+1} = n_{j+1}$, then $|B_{j,j+1}| = |\mathcal{R}_j|$. Let

$$\Delta_j = \begin{cases} L_{j-1,j} & \text{if } k_{j-1} + k_j = n_j, \\ B_{j,j+1} & \text{if } k_j + k_{j+1} = n_{j+1}, \\ L_{j-1,j} \text{ or } B_{j,j+1} & \text{if } k_{j-1} + k_j = n_j \text{ and } k_j + k_{j+1} = n_{j+1}. \end{cases}$$

We have $|\Delta_j| = |\mathcal{R}_j|$, and the rows of J indexed by elements in \mathcal{R}_j are identical to the rows of J indexed by Δ_j . It is easily seen that the rows of J_x indexed by the codim V elements in the sets

$$\mathcal{R}_1, \dots, \mathcal{R}_{j-1}, \Delta_j, \mathcal{R}_{j+1}, \dots, \mathcal{R}_h, K_{1,2}, \dots, K_{h-1,h}$$

are linearly independent, and hence x is smooth.

LEMMA 2.8. *Let $x = (A_1, \dots, A_h)$ be a point of V such that if $\text{rank } A_j < k_j$ for some $1 \leq j \leq h$, then*

1. *either $1 \leq j-1 \leq h$ and $\text{rank } A_{j-1} = k_{j-1}$, or $1 \leq j+1 \leq h$ and $\text{rank } A_{j+1} = k_{j+1}$;*
2. *either $k_{j-1} + k_j = n_j$, or $k_j + k_{j+1} = n_{j+1}$ (here $k_0 = k_{h+1} = 0$).*

Then x is a smooth point of V .

Proof. Let Δ_j be as in Remark 2.7. Then the rows of J_x indexed by the codim V elements in the sets

\mathcal{R}_i , with i such that $\text{rank } A_i = k_i$,

Δ_j , with j such that $\text{rank } A_j < k_j$,

$K_{1,2}, \dots, K_{h-1,h}$

are linearly independent. Therefore x is a smooth point of V . ■

Remark 2.9. An alternative proof of Lemma 2.8 can be obtained as follows. For $1 \leq j \leq h$ with $\text{rank } A_j < k_j$, let Γ_j be as in the proof of Lemma 2.6. It is easily seen that the columns of J_x that are indexed by the elements in the disjoint sets

Y_i , with i such that $\text{rank } A_i = k_i$,

$Q_{i,i+1}^+$ with i such that $\text{rank } A_i = k_i$, $\text{rank } A_{i+1} = t_{i+1}$,

Γ_j , with j such that $\text{rank } A_j < k_j$

are linearly independent. Since the number of these columns is codim V , we conclude that $\text{rank } J_x = \text{codim } V$, and hence x is smooth.

LEMMA 2.10. *Let $x = (A_1, \dots, A_h)$ be a point of V such that $\text{rank } A_j < k_j$ and $\text{rank } A_{j+1} < k_{j+1}$ for some j , $1 \leq j \leq h-1$. Then x is a singular point of V .*

Proof. Let $\text{rank } A_i = t_i$, with $t_j < k_j$, $t_{j+1} < k_{j+1}$. Without loss of generality, we may assume that

$$A_i = \begin{pmatrix} 0 & 0 \\ N_i & 0 \end{pmatrix}, \quad \text{for } 1 \leq i \leq h,$$

where N_i is the $t_i \times t_i$ identity matrix.

The nonzero columns of J_x are among the nonzero columns of J . Let T be the indices of a maximal set of linearly independent columns of J_x indexed by entries in $X_1, \dots, X_{j-1}, X_{j+2}, \dots, X_h$. Note that a column of J indexed by an entry in $X_1, \dots, X_{j-1}, X_{j+2}, \dots, X_h$ is either identical to some column indexed by an entry in $X_j \cup X_{j+1}$ or linearly independent of the columns indexed by entries in $X_j \cup X_{j+1}$. Thus there exists a set $S \subset X_j \cup X_{j+1}$ such that the columns of J_x indexed by elements of $S \cup T$ give a maximal set of linearly independent columns of J_x , and we have $\text{rank } J_x = |S| + |T|$.

As noted in Remark 2.4, there are precisely $\text{codim } V - (|Y_j| + |Y_{j+1}| + |Q_{j,j+1}^\pm|)$ linearly independent columns of J indexed by entries in $X_1, \dots, X_{j-1}, X_{j+2}, \dots, X_h$. Therefore $|T| \leq \text{codim } V - (|Y_j| + |Y_{j+1}| + |Q_{j,j+1}^\pm|)$.

Now we show that $|S| < |Y_j| + |Y_{j+1}| + |Q_{j,j+1}^\pm|$. Each column indexed by an element of S contains precisely one nonzero entry in a row indexed by an entry of $X_j X_{j-1}, X_{j+1} X_j$, or $X_{j+2} X_{j+1}$ (since $\text{rank } A_j < k_j$ and $\text{rank } A_{j+1} < k_{j+1}$). The columns indexed by entries in $Q_{j-1,j}^-$ and $Q_{j+1,j+2}^+$ are identical to columns in T (see Remark 2.4). Therefore $S \cap Q_{j-1,j}^- = \emptyset$, $S \cap Q_{j+1,j+2}^+ = \emptyset$, and $S \subset Y_j \cup Y_{j+1} \cup Q_{j,j+1}^+ \cup Q_{j,j+1}^-$. The set of columns indexed by entries in $Q_{j,j+1}^+$ is the same as the set of columns indexed by entries in $Q_{j,j+1}^-$, so we may assume that $S \cap Q_{j,j+1}^- = \emptyset$. Thus we have $|S| = |S \cap Y_j| + |S \cap Y_{j+1}| + |S \cap Q_{j,j+1}^+|$. Obviously, $|S \cap Y_j| \leq |Y_j|$ and $|S \cap Y_{j+1}| \leq |Y_{j+1}|$ (note that one can have equalities here). On the other hand, we have $|S \cap Q_{j,j+1}^+| \leq t_j t_{j+1} < k_j k_{j+1} = |Q_{j,j+1}^\pm|$. Consequently, $|S| < |Y_j| + |Y_{j+1}| + |Q_{j,j+1}^\pm|$.

Thus $\text{rank } J_x < \text{codim } V$, which shows that x is a singular point of V .

Next we describe the singular locus of V .

For $1 \leq i \leq h$, let V_i denote the set of all points (A_1, \dots, A_h) in V such that $\text{rank } A_i < k_i$, i.e.,

$$V_i = V(k_1, \dots, k_{i-1}, k_i - 1, k_{i+1}, \dots, k_h).$$

For $1 \leq j \leq h - 1$, let $V_{j,j+1}$ denote the set of all points (A_1, \dots, A_h) in V such that $\text{rank } A_j < k_j$ and $\text{rank } A_{j+1} < k_{j+1}$, i.e.,

$$V_{j,j+1} = V(k_1, \dots, k_{j-1}, k_j - 1, k_{j+1} - 1, k_{j+2}, \dots, k_h).$$

Note that $V_{j,j+1} = V_j \cap V_{j+1}$.

THEOREM 2.11. *The irreducible components of $\text{Sing } V$ are the subvarieties of V of the form V_i , with $i \in \Omega$, and $V_{j,j+1}$, with $j \notin \Omega$, $j + 1 \notin \Omega$, where Ω is the set of all $1 \leq i \leq h$ such that $k_{i-1} + k_i < n_i$ and $k_i + k_{i+1} < n_{i+1}$.*

Proof. Note that $\text{Sing } V$ is a union of G_n -orbit closures, i.e., subvarieties of V of the form $V(\mathbf{r})$, with $\mathbf{r} \in \mathcal{K}_n$.

Next we determine for which $\mathbf{r} = (r_1, \dots, r_h) \in \mathcal{K}_n$ the subvariety $V(\mathbf{r})$ is singular. To check that $V(r_1, \dots, r_h)$ is singular, it is enough to check that the point $x = (A_1, \dots, A_h)$ with $\text{rank } A_i = r_i$ is singular.

If there is no i with $r_i < k_i$, by Lemma 2.1 $V(\mathbf{r})$ is not singular.

If there is precisely one index i with $r_i < k_i$, by Lemma 2.6 the subvariety $V(\mathbf{r}) = V_i$ is singular if and only if $i \in \Omega$.

Next, we analyze the subvarieties $V(\mathbf{r})$ with $r_i < k_i$ for more than one index i .

If there is some j with $r_j < k_j$ and $r_{j+1} < k_{j+1}$, then $V(\mathbf{r}) \subset V_{j,j+1}$. By Lemma 2.10 $V_{j,j+1}$ is singular, and therefore $V(\mathbf{r})$ is singular.

If for each j we have either $r_j = k_j$ or $r_{j+1} = k_{j+1}$, then by Lemma 2.8 $V(\mathbf{r})$ is singular only if $i \in \Omega$ for some i with $r_i < k_i$. But in this case $V(\mathbf{r}) \subset V_i$, and since V_i is singular, $V(\mathbf{r})$ is also singular.

Thus we have determined all the subvarieties of V of the form $V(\mathbf{r})$ with $\mathbf{r} \in \mathcal{K}_n$, $\mathbf{r} \leq \mathbf{k}$, that are contained in $\text{Sing } V$. By taking the maximal such subvarieties (with respect to set inclusion), we obtain the required description for the irreducible components of $\text{Sing } V$. ■

As a consequence, we obtain the following result that describes the singular loci of the irreducible components of the variety of complexes:

THEOREM 2.12. *If V is an irreducible component of the variety of complexes \mathcal{C} , then the irreducible components of $\text{Sing } V$ are $V_{1,2}, \dots, V_{h-1,h}$.*

Proof. In this case the set Ω is empty (by Theorem 1.2), and the result follows immediately from Theorem 2.11. ■

REFERENCES

1. S. Abeasis and A. Del Fra, Degenerations for the representations of a quiver of type A_m , *J. Algebra* **93** (1985), 376–412.
2. C. De Concini and E. Strickland, On the variety of complexes, *Adv. Math.* **41** (1981), 57–77.
3. V. Lakshmibai, Singular loci of varieties of complexes, preprint, 1998.
4. V. Lakshmibai and P. Magyar, Degeneracy schemes, Schubert varieties and quiver varieties, *Internat. Math. Res. Notices* **2** (1998), 627–640.
5. C. Musili and C. S. Seshadri, Schubert varieties and the variety of complexes: Arithmetic and geometry, Vol. II, *Progr. Math.* **36** (1983), 329–359.